

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Letter to the Editor

Letter to the Editor: Validity and reliability concerns associated with cardiopulmonary exercise testing young people with cystic fibrosis. Response to: Statement on Exercise Testing in Cystic Fibrosis (Hebestreit *et al.*, 2015 *Respiration* 90(4):332-51)

Zoe L. Saynor^{a*}, Dr. Alan R. Barker^b, Dr. Patrick J. Oades^c, Owen W. Tomlinson^b, Prof. Craig Anthony Williams^b

Word count: 920 words

Short Title: Validity and reliability concerns associated with CPET in CF

26

27

28

29

30 ^a Department of Sport and Exercise Science, University of Portsmouth, Portsmouth, UK.

31

32 ^b Children's Health and Exercise Research Centre, Sport and Health Sciences, University of
33 Exeter, Exeter, Devon, UK.

34

35 ^c Paediatric unit, Royal Devon and Exeter NHS Foundation Trust, Exeter, Devon, UK.

36

37 *Correspondence to: Z.L. Saynor, Department of Sport and Exercise Science, Spinnaker
38 Building, Cambridge Road, University of Portsmouth, Portsmouth, Hampshire, UK.

39 Tel: +44 (0)2392 843080

40 Email: zoe.saynor@port.ac.uk

41 The recent statement by Hebestreit and colleagues [1] on behalf of the European Cystic
42 Fibrosis Society (ECFS) Exercise Working Group and endorsed by the European Respiratory
43 Society, should be commended for their efforts to establish consensus regarding exercise
44 testing for young people with CF. Exercise testing is a valuable investigative tool within both
45 the clinical management and scientific investigation of children and adolescents with CF and
46 this document provides an international standpoint regarding the importance of
47 comprehensive cardiopulmonary exercise testing (CPET) within the clinical management of
48 this patient group. However, it is our view that the authors have missed an opportunity to
49 provide a contemporary overview of the CPET ‘toolkit’ currently available.

50 The authors state that this document will *‘describe the current best practice*
51 *recommendations for conducting exercise tests in patients with CF’* and *‘summarises the*
52 *information available on specific test protocols and outcome parameters (Page 2)’*. The
53 authors recommend the Godfrey protocol [2] when using the cycle ergometer, with measures
54 of arterial oxygen saturation and, when possible, pulmonary gas exchange and ventilation.
55 Whilst this does represent progress from the routinely used shuttle and step tests, the authors
56 failed to acknowledge several limitations inherent to the Godfrey protocol and the
57 recommended use of criteria to verify a maximal test. This is surprising, given that the ECFS
58 Clinical Trials Network Standardisation Committee recently called for research assessing the
59 validity, reproducibility and feasibility of outcome measures utilised in the assessment of
60 patients with CF and the most appropriate exercise test for paediatric patients [3].

61 The authors rightfully acknowledge that an issue with shuttle and step tests is that it can be
62 difficult to determine whether a maximal effort was made. However, they then state that *‘the*
63 *Godfrey protocol provides valid information for all CF relevant indications for an exercise*
64 *test’*. The authors recommend that since not all individuals display the tradition verification
65 criterion of a plateau in oxygen uptake (VO_2) upon exhaustion, at least one of the following

should be used to confirm a maximal effort: the patient achieves a predicted $\text{VO}_{2\text{peak}}$ or peak power output (W_{peak}); the patient reaches maximal heart rate, peak ventilation approaches maximal voluntary ventilation, respiratory exchange ratio (RER) is > 1.03 , exertion is 9-10 on the 0-10 scale or ≥ 17 on a 7-20 scale. However, our research group recently demonstrated that the use of secondary criteria to confirm a maximal effort (e.g. $\text{RER} > 1.00$ or 1.10 , [blood lactate] $\geq 6 \text{ mmol} \cdot \text{L}^{-1}$), as recommended by Hebestreit *et al.* [1], are invalid and can drastically underreport maximal $\text{VO}_{2\text{max}}$ in some young people with CF [4], a finding consistent with healthy children and adolescents [5]. Accepting submaximal or rejecting ‘true’ maximal values could distort the clinical application and interpretation of CPET in young people with CF.

We have, however, demonstrated that a procedure termed the ‘supramaximal verification phase’ (S_{max}), in which an exhaustive ramp incremental test precedes an exhaustive individualised constant work rate test at an intensity above W_{peak} , can confirm whether a ‘true’ measure of $\text{VO}_{2\text{max}}$ has been obtained, which is fundamental to the utility of this outcome parameter in CF. This finding is in line with data in healthy adults [6-14], children [5] and other paediatric clinical groups [15]. Although the authors present information regarding ‘*was the test maximal?*’, they failed to mention this published evidence and presented inaccurate verification criteria as *best* CPET practice for young people with CF, which should be approached with caution. This statement also provides a summary of the reliability of exercise tests for young people with CF, however again recent evidence has been ignored. We recently reported both the short- and medium-term reproducibility of a valid CPET protocol for young people with CF [16], which was shown to reduce the error of measurement when compared with an isolated incremental CPET to derived $\text{VO}_{2\text{peak}}$ [16].

Whilst the focus of this letter addresses validity and reproducibility issues with the Godfrey protocol, other potential issues to consider are: ‘step’ increases in work rate derived exclusively from stature can result in insufficient test durations of ≤ 4 minutes in young people with CF [17], which limits our ability to characterise the progressive increase in VO_2 during exercise and determine submaximal measures of aerobic fitness (e.g. the gas exchange threshold or VO_2 mean response time) which, as highlighted in this consensus statement, may provide better predictors of mortality in adolescents with CF [18]. In accordance with others [19], we therefore recommend a ramp incremental exercise test, which aims to reach volitional exhaustion in 8-12 minutes [20], followed by S_{max} verification of maximal CPET parameters. Not only has this testing protocol been demonstrated as safe and feasible in young people with CF in a research setting, it is also now used as part of patients’ annual clinical review with UK based CF clinics in Exeter, Southampton and Portsmouth, demonstrating the feasibility of its clinical implementation. The CF-specific linear regression model to predict W_{peak} and calculate individualised workload increments to reach volitional exhaustion in ~ 10 minutes developed by Hulzebos and colleagues should help prevent short test durations [21].

Whilst it is recognised that there are not currently any large scale studies directly comparing exercise testing protocols, we feel the authors could have provided a more contemporary overview of the evidence concerning the validity and reproducibility of CPET protocols available for use in young people with CF. If the clinical utility of CPET to provide a comprehensive evaluation of physiological (dys)function and stratify patients with CF is to be realised, these important practical considerations much be acknowledged.

REFERENCES

1. Hebestreit H, Arets HG, Aurora P, Cerny F, Hulzebos EH, Karila C, Lands LC, Lowman JD, Swisher A, Urquart DS; European Cystic Fibrosis Exercise Working Group: Statement on exercise testing in cystic fibrosis. *Respiration* 2015;90(4):332-51.
2. Godfrey S, [Davies CT](#), Wozniak E, [Barnes CA](#): Cardio-respiratory response to exercise in normal children. *Clin Sci* 1971;40(5):419-31.
3. Bradley JM, Madge S, Morton AM, Quittner AL, Elborn JS: Cystic fibrosis research in allied health and nursing professions. *J Cyst Fibros* 2012;11(5):387-392.
4. Saynor ZL, Barker AR, Oades PJ, Williams CA: A protocol to determine valid $\text{VO}_{2\text{max}}$ in young cystic fibrosis patients. *J Sci Med Sport* 2013;16(6):539-544.
5. Barker AR, Williams CA, Jones AM, Armstrong N: Establishing maximal oxygen uptake in young people during a ramp cycle test to exhaustion. *Br J Sports Med* 2011;45(6):498-503.
6. Day JR, Rossiter HB, Coats EM, Skasick A, Whipp BJ: The maximally attainable VO_2 during exercise in humans: the peak vs. maximum issue. *J Appl Physiol* 2003;95(5):1901-1907.
7. [Midgley AW](#), [McNaughton LR](#), [Carroll S](#): Verification phase as a useful tool in the determination of the maximal oxygen uptake of distance runners. *Appl Physiol Nutr Metab* 2006;31(5):541-8.
8. Rossiter HB, Kowalchuk JM, Whipp BJ: A test to establish maximum O_2 uptake despite no plateau in the O_2 uptake response to ramp incremental exercise. *J Appl Physiol* 2006;100(3):764-770.
9. [Midgley AW](#), [McNaughton LR](#), [Polman R](#), [Marchant D](#): Criteria for determination of maximal oxygen uptake: a brief critique and recommendations for future research. *Sports Med* 2007;37(12):1019-28.
10. [Hawkins MN](#), Raven PB, Snell PG, Stray-Gundersen J, Levine BD: Maximal oxygen uptake as a parametric measure of cardiorespiratory capacity. *Med Sci Sports Exerc* 2007; 39(1):103-7.
11. Foster C, Kuffel E, Bradley N, Battista RA, Wright G, Porcari JP, Lucia A, deKoning JJ: $\text{VO}_{2\text{max}}$ during successive maximal efforts. *Eur J Appl Physiol* 2007;102(1):67-72.
12. Poole DC, Wilkerson DP, Jones AM: Validity for establishing maximal O_2 uptake

during ramp exercise tests. Eur J Appl Physiol 2008;102(4):403-410.

13. Midgley AW, Carroll S. Emergence of the verification phase procedure for confirming 'true' VO₂max. Scand J Med Sci Sports 2009;19(3), 313-22.

14. Scharhag-Rosenberger F, Carlsohn A, Cassel M, Mayer F, Scharhag J: How to test maximal oxygen uptake: a study on timing and testing procedure of a supramaximal verification test. Appl Physiol Nutr Metab 2011;36(1):153-160.

15. de Groot JF, Takken T, de Graaf S, Gooskens RH, Helders PJ, Vanhees L: Treadmill testing of children who have spina bifida and are ambulatory: does peak oxygen uptake reflect maximum oxygen uptake? Phys Ther 2009;89(7):679-687.

16. Saynor ZL, Barker AR, Oades PJ, Williams CA: Reproducibility of maximal cardiopulmonary exercise testing for young cystic fibrosis patients. J Cyst Fibros 2013;12(6):644-650.

17. Kent L, O'Neill B, Davidson G, Nevill A, Murray J, Reid A, Elborn JS, Bradley JM: Cycle ergometer tests in children with cystic fibrosis: reliability and feasibility. Pediatr Pulmonol 2012;47(12):1226-1234.

18. [Hulzebos EH](#), [Bomhof-Roordink H](#), [van de Weert-van Leeuwen PB](#), [Twisk JW](#), [Arets HG](#), [van der Ent CK](#), [Takken T](#): Prediction of mortality in adolescents with cystic fibrosis. Med Sci Sports Exerc 2014;46(11):2047-52.

19. Bongers BC, van Brussel M, Hulzebos HJ, Takken T: Paediatric exercise testing in clinics and classrooms: A comparative review of different assessments. OA Epidemiology 2013;1(2):14.

20. Williams CA, Saynor ZL, Tomlinson OW, Barker AR: Cystic fibrosis and physiological responses to exercise. Expert Rev Respir Med 2014;8(6):751-752.

21. Hulzebos HJ, Werkman MS, van Brussel M, Takken T: Towards an individualized protocol for workload increments in cardiopulmonary exercise testing in children and adolescents with cystic fibrosis. J Cyst Fibros 2012;11(6):550-554.